

Oxidized Thioredoxin Reductase Assay Kit (Spectrophotometry)

Description

Thioredoxin reductase (Oxidized Thioredoxin Reductase, TrxR) is an NADPH-dependent, FAD-containing dimeric selenoenzyme that belongs to the pyridine nucleotide–disulfide oxidoreductase family. Together with thioredoxin and NADPH, it forms the thioredoxin system.

TrxR exhibits activity similar to glutathione reductase (GR), catalyzing the reduction of GSSG to GSH, and is one of the key enzymes in the glutathione redox cycle.

Detection Principle

Thioredoxin reductase (TrxR) catalyzes the reduction of DTNB by NADPH to generate TNB and NADP⁺. TNB exhibits a characteristic absorption peak at 412 nm. However, reduced glutathione (GSH) can also react with DTNB to produce TNB. Therefore, 2-vinylpyridine is used to inhibit endogenous reduced glutathione in the sample. The activity of TrxR can then be calculated by measuring the rate of increase in TNB absorbance at 412 nm.

Packing

Taking 50T/48S packing for example:

Components	Packing	Storage
CB0136S-A	100 mL x 1	4 °C
CB0136S-B	1 vial (powder) x 1	Store at 4 °C protected from light; before use, dissolve by adding 5 mL of 1× PBS buffer (use within 3 days).
CB0136S-C	1 vial (powder) x 1	Store at 4 °C; before use, dissolve by adding 5 mL of distilled water (use within 3 days).
CB0136S-D	30 μL×1	Store at -20 °C; before use, dilute 10-fold with absolute ethanol according to the number of samples.

Prior to the formal determination, a preliminary assay should be conducted using 2-3 samples with large expected differences.

Instructions

I. Preparation of Lab Instruments

Visible spectrophotometer, 1 mL glass cuvette, water bath, balance, centrifuge, adjustable pipette, mortar, distilled water, 1× PBS buffer.

II. Extraction of Crude Enzyme

1. Tissues:

Homogenize on ice at a ratio of tissue mass (g) : CB0136S-A volume (mL) = 1 : 5–10

(Recommended: weigh ~0.1 g tissue and add 1 mL CB0136S-A).

Centrifuge at 8000 × g, 4 °C for 10 min. Collect the supernatant and keep on ice for analysis.

2. Bacteria / cells:

Use a ratio of cell number (×10⁴ cells) : CB0136S-A volume (mL) = 500–1000 : 1

(Recommended: add 1 mL CB0136S-A to 5 × 10⁶ cells).

Lyse cells by ultrasonication on ice (300 W; 3 s on / 7 s off; total 3 min).

Then centrifuge at 8000 × g, 4 °C for 10 min. Collect the supernatant and keep on ice for analysis.

3. Serum and other liquid samples: Measure directly.

III. Assay Procedure

1. Preheat the spectrophotometer for 30 minutes, set the wavelength to 412 nm, and zero the instrument using distilled water.
2. Preheat CB0136S-A at 25 °C (for general species) or 37 °C (for mammals) for 30 minutes. Before measurement, mix the sample supernatant with CB0136S-D at a volume ratio of 50:1 (i.e., add 2 μL of CB0136S-D to 100 μL of supernatant). Incubate in a 37 °C water bath for 30 minutes, then place on ice.
3. Add the following reagents

	Blank Tube (μL)	Sample Tube (μL)
CB0136S-B	100	
CB0136S-C	100	
CB0136S-A	800	
After rapid mixing, measure the absorbance at 412 nm at 10 s. Then incubate in a 37 °C water bath for 5 min, quickly remove and measure the absorbance at 412 nm again, recorded as A1 and A2. Calculate the blank tube ΔA as ΔA _{blank} = A2 – A1.		
CB0136S-B		100
CB0136S-C		100
CB0136S-A		700
supernatant		100
After rapid mixing, measure the absorbance at 412 nm at 10 s. Then incubate in a 37 °C water bath for 5 min, promptly remove and measure the absorbance at 412 nm again. Record the two readings as A3 and A4, respectively. Calculate the absorbance change for the assay tube as ΔA = A4 – A3.		

Note: Blank tubes only need to be measured 1–2 times.

IV. Calculation of TrxR Activity

1. Based on protein concentration

At 25 °C or 37 °C, the amount of enzyme that catalyzes the reduction of 1 nmol of DTNB per minute per milligram of protein is defined as one unit of enzyme activity.

$$\text{TrxR (U/mg prot)} = (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div \epsilon \div d \times 109 \times V1 \div (C_{\text{pr}} \times V2) \div T = 147 \times (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div C_{\text{pr}}$$

2. Based on sample mass

At 25 °C or 37 °C, one enzyme activity unit is defined as the amount of enzyme that catalyzes the reduction of 1 nmol of DTNB per minute per gram of sample.

$$\text{TrxR (U/g)} = (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div \epsilon \div d \times 109 \times V1 \div (W \times V2 \div V3) \div T = 147 \times (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div W$$

3. Based on bacterial or cell density

At 25 °C or 37 °C, one enzyme activity unit is defined as the amount of enzyme that catalyzes the reduction of 1 nmol of DTNB per minute per 10⁴ cells.

$$\text{TrxR (U/104 cell)} = (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div \epsilon \div d \times 109 \times V1 \div (\text{cell number} \times V2 \div V3) \div T = 147 \times (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div \text{cell number}$$

4. Based on Liquid Volume

At 25 °C or 37 °C, one enzyme activity unit is defined as the amount of enzyme that catalyzes the reduction of 1 nmol of DTNB per minute per milliliter of liquid.

$$\text{TrxR (U/mL)} = (\Delta A_{\text{sample}} - \Delta A_{\text{blank}}) \div \epsilon \div d \times 109 \times V1 \div V2 \div T = 147 \times (\Delta A_{\text{sample}} - \Delta A_{\text{blank}})$$

Note:

ε: Molar extinction coefficient of TNB at 412 nm, 1.36 × 10⁴ L/mol/cm; d: Optical path length of the cuvette, 1 cm

V1: Total volume of the reaction system (L), 1000 μL = 1 × 10⁻³ L; C_{pr}: Protein concentration of the supernatant (mg/mL), to be determined separately

W: Sample mass; V2: Volume of supernatant added to the reaction system (mL), 100 μ L = 0.1 mL

V3: Volume of extraction buffer, 1 mL; T: Reaction time (min), 5 min

Cell number: Expressed in units of 10^4 cells (ten thousand cells); 10^9 : Unit conversion factor, 1 mol = 10^9 nmol





Precautions

1. Before formal measurement, perform a preliminary test using 1–2 samples to ensure that the absorbance shows a linear change within 5 minutes. For TrxR activity assays of mammalian tissues and blood products, samples generally need to be diluted approximately 5-fold with distilled water. All procedures during the assay should be carried out rapidly.
2. Since CB0136S-A contains a certain concentration of protein (approximately 0.1 mg/mL), the protein contributed by the extraction buffer itself must be subtracted when determining the protein concentration of the samples.
3. The product is for R&D use only, not for diagnostic procedures, food, drug, household or other uses.
4. Please wear a lab coat and disposable gloves.

TargetMol US

 sales@targetmol.com  (781) 999-4286  www.targetmol.com
 34 Washington Street, Suite 220, Wellesley Hills, MA 02481

TargetMol EU

 sales@targetmol.com  +43(0)676/786025  www.targetmol.com
 Hafenstraße 47-51, 4020 Linz, Austria



LinkedIn



Facebook



PDF Documents